

## The systematic search for $z \geq 5$ active galactic nuclei in the *Chandra* Deep Field South

Anna K. Weigel<sup>1</sup>, Kevin Schawinski<sup>1</sup>, Ezequiel Treister<sup>2</sup>, C. Megan Urry<sup>3</sup>,  
Michael Koss<sup>1</sup>, Benny Trakhtenbrot<sup>1</sup>

<sup>1</sup> *ETH Zurich, Zurich, Switzerland*

<sup>2</sup> *Universidad de Concepción, Concepción, Chile*

<sup>3</sup> *Yale University, New Haven, United States*

### Abstract

We investigate early black hole growth through the methodical search for  $z \geq 5$  AGN in the *Chandra* Deep Field South. We base our search on the *Chandra* 4-Ms data which at  $z \sim 5$  makes us sensitive to luminosities as low as  $\sim 10^{42} \text{ erg s}^{-1}$  ( $\sim 10^{43} \text{ erg s}^{-1}$ ) in the soft (hard) band. The deep X-ray data should allow us to detect Compton-thin AGN with  $M_{\text{BH}} > 10^7 M_{\odot}$  accreting at Eddington ratios  $> 0.1$ . The field contains over 600  $z \sim 5$  Lyman Break Galaxies. These high-redshift galaxies are the progenitors of massive, local galaxies and based on lower redshift relations we would expect  $\sim 20$  of them to host AGN. We combine the *Chandra* data with optical GOODS/ACS and infrared CANDELS/WFC3 and *Spitzer*/IRAC data. After excluding clear low-redshift sources and objects with insufficient filter coverage, our main sample consists of 58 high-redshift candidates. We use a range of redshift estimators including a photo- $z$  code, stacking, colour criteria and the Lyman Break Technique. We also use the X-ray Hardness Ratio as additional information. The final  $z \geq 5$  candidates that remain after we combine our redshift tests, are likely to be low-redshift interlopers. We thus conclude that, contrary to our expectation of finding at least a few high-redshift AGN, the field does not contain any convincing  $z \geq 5$  AGN candidates. Our results place interesting constraints on early black hole growth and we discuss a range of possible explanations.